



**AG-NUTRI
PTY. LTD.**

LIG-CALCIUM + BORON

Guaranteed Analysis

Calcium 6% as calcium lignosulphonate chelate
Boron 2% as ammonium polyborate complex

The proven calcium formulation
for superior crop quality

▶ Fully chelated liquid formulated to correct deficiencies in fast growing crops

▶ Binding properties provide more efficient utilization of nutrients

▶ Contains boron to assist soil calcium uptake

▶ Very safe & gentle on crops (contains no chloride or sodium-based chelating agents)

▶ Excellent nutrient & chemical compatibility

“We don't just sell a product, we prescribe a program”



Apples - Apply 5 L/ha prior to flowering & 5-6 sprays @ 7.5 L/ha commencing 10 days after petal fall up until 4 weeks prior to harvest.



Avocado - Apply LIGNO-BORON @ 4 L/ha 21 days prior to flowering & every 21 days from flower bud development to fruit set. From fruit set to harvest apply LIG-CALCIUM + BORON @ 5 L/ha every 21 days & LIGNO-BORON @ 4 L/ha post harvest.

Brassica - Apply LIG-CALCIUM + BORON @ 5 L/ha prior to head/heart formation & repeat every 10-14 days until harvest.

Capsicum - Apply 7.5 L/ha over the leaf & up the drip just prior to bud development & repeat every 14 days.

Carrots - Apply 7.5 L/ha when plants are 10 cm tall plus 2 more sprays during mid-growth stage.

Citrus - Apply 4-6 sprays @ 5 L/ha commencing at flowering every 10-14 days. Include 3 L/ha VIGOR-LIG PLUS as a binder and replacement for urea.

Cucurbits - Apply 5 L/ha at 1st flower & repeat every 14 days during fruit set to harvest with Calcium + Boron PLEX @ 7.5 L/ha up the drip.



Mango - Fertigate 21 days prior to flowering Calcium + Boron PLEX @ 10 L/ha & repeat every 14 days from fruit set to harvest. Apply LIGNO-BORON @ 4 L/ha every 14 days from bud initiation to fruit set. From fruit set to harvest apply LIG-CALCIUM + BORON @ 5 L/ha every 14 days & LIGNO-BORON @ 4 L/ha post harvest.

Olives - Prior to flowering apply 2 sprays @ 5 L/ha (or LIGNO-BORON @ 2.5 L/ha) & from fruit set to harvest repeat every 4 weeks. On young trees apply 5 L/500 L water as required.



Onions - Apply 2-3 sprays @ 7.5 L/ha commencing at mid-growth every 14-21 days.

Potatoes - Apply LIG-CALCIUM + BORON @ 7.5 L/ha after tuber set & repeat 2-3 weeks later with LIG-K + BORON @ 10 L/ha.

Stonefruit - 4-6 sprays @ 5 L/ha commencing 7-10 days after petal fall & repeat every 7-14 days.



Strawberries - Apply 7.5 L/ha as a foliar spray plus Calcium + boron PLEX @ 10 L/ha up the drip every 14 days during the growing season. For berry sizing also include LIG-K + BORON @ 5-10 L/ha.

Tomatoes - Apply 7.5 L/ha at 1st flower & repeat every 21 days until fruit is formed. For hydroponic tomatoes apply 1 L/100 L water every 14 days.

Vegetables - Apply 5 L/ha prior to flowering, at mid-growth or at head/heart formation & repeat every 14-21 days until harvest.

Vines - Apply 5 L/ha prior to flowering & at fruit set. Repeat at bunch closure, Veraison & pre-harvest in table grapes & sensitive wine grape varieties.

Also talk to us about our liquid calcium nitrate equivalent - CR CAL-NIT + Boron.



CONTACTS

National Manager
Steven Brooks (Dip. Ag. Sc.)
0418 520 744
steve@sjbagnutri.com.au

South Australia, Sunraysia & WA
Nathan Smith (B. Ag.)
0408 633 661
nathan@sjbagnutri.com.au

Queensland & NT
Mark Bingham (B. Ag. Sc.)
0439 520 744
mark@sjbagnutri.com.au

NSW & Technical Support
Leigh Teitz (B. App. Sc. Ag.)
0407 040 820 Fax: 0358 621 279
leigh@sjbagnutri.com.au