



**AG~NUTRI
PTY. LTD.**

LIG-K + BORON

Potassium 10% + Boron 2%

Potassium is required for:

- sugar production (increases baume in vines & total solids in tomatoes)
- fruit filling & size (eg. strawberries, grapes & carrots)
- crop resistance to disease, moisture stress & frost

Boron is required for:

- calcium uptake
- sugar transport & fruit filling
- cell wall strength & skin quality (prevents flaky skins in potatoes)



The advantages of using LIG-K + BORON

- chelated formulation rapidly absorbed & translocated to developing fruit
- compatible with other chemical & nutrient sprays
- very safe on fruit (ie. no marking on table grapes)
- produces healthier, more robust plants
- increases fruit size & quality



Application of LIG-K + BORON

Carrots - 2-3 sprays @ 10 L/ha starting at mid-growth stage.

Onions - 2-3 sprays @ 10 L/ha starting at bulb filling.

Potatoes - 2-3 sprays @ 10 L/ha commencing at tuber set

Strawberries - 5-10 L/ha every 14 days during fruiting

Tomatoes - 2-3 sprays @ 10 L/ha starting at fruit set

Vines - 5 L/ha prior to flowering, at fruit set & bunch closure. For table grapes apply 4-5 sprays @ 5 L/ha between fruit set to harvest.